

FLIGHT PROGRAM TO STUDY SPACECRAFT/ATMOSPHERIC INTERACTIONS

A. L. Broadfoot

**University of Arizona
Lunar and Planetary Laboratory
Tucson, Arizona 85721**

November 2000

Final Report

20020528 039

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.



**AIR FORCE RESEARCH LABORATORY
Space Vehicles Directorate
29 Randolph Road
AIR FORCE MATERIEL COMMAND
HANSCOM AFB MA 01731-3010**

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE November 2000		3. REPORT TYPE AND DATES COVERED FINAL REPORT (Oct 91 - Sep 99)
4. TITLE AND SUBTITLE FLIGHT PROGRAM TO STUDY SPACECRAFT/ATMOSPHERIC INTERACTION			5. FUNDING NUMBERS PE: 63217F PR: S321 TA: 42 WU: AD Contract #: F19628-91-K-0041	
6. AUTHOR(S) A. L. Broadfoot				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Lunar and Planetary Laboratory University of Arizona Tucson, Arizona 85721			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory 29 Randolph Road Hanscom AFB MA 01731-3010			10. SPONSORING/MONITORING AGENCY REPORT NUMBER AFRL-VS-TR-2001-1583	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This effort was in preparation for a continuing study of spacecraft/atmosphere interactions.				
14. SUBJECT TERMS Space flight Atmosphere interactions Spacecraft			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT SAR	

Arizona Airglow Experiment (GLO): Flight Program to Study Spacecraft/Atmosphere Interactions

The Arizona Airglow Experiment (GLO) used a Space Shuttle-borne spectrograph and imager suite to study the atmosphere from 60 to 400 km and the *in situ* interactions between the Space Shuttle and the ionosphere. Published reports include data on thruster interactions with the ionosphere, nascent dayglow and night airglow of the atmosphere, and metal atom and ion emissions in the thermosphere.

Two essentially identical GLO instruments were constructed and flew a total of six times on board 5 separate shuttle flights, with real-time ground control from NASA Goddard Space Flight Center in Greenbelt, MD. Each shuttle flight provided a unique set of orbital parameters affecting the ionospheric observations, so a much greater dataset was obtained than would have been possible if the flights had been identical. The experiment returned over 50 gigabytes of hyperspectral data, where spectral and altitude dimensions are recorded directly from the detectors, and distance along flight track is accumulated over time for each orbit.

Flight parameters for each mission are summarized in Table 1, showing the shuttle orbit and atmospheric observations for each GLO. The shuttle orbit alignment is shown in Figure 1. The first GLO flight on STS-53 closely followed the terminator, and provided excellent viewing of the dawn and dusk atmospheres. The remaining GLOs flew orbits that crossed the terminators sharply, providing various combinations of time-of-day and observable latitude. GLO-5 and GLO-6 flew together on STS-85 (see Figure 2), and their views were arranged to provide (i) greater dayglow tangent height coverage than that afforded by a single machine, (ii) different angles through the night airglow atmosphere to provide multiple height resolutions, or (iii) near-stereo viewing to aid in 3-dimensional projections of the atmosphere. For the latter, one GLO viewed typically ahead of the shuttle track and the other viewed behind the track to scan the same patch of ionosphere from two angles with only a short time difference. The STS-85 end-of-mission summary is included as Appendix I.

Table 1. GLO Flight Parameters

GLO-	1	2	3	4	5 & 6
STS-	53	63	69	74	85
Orbital Inclination	57°	51°	28°	51°	57°
Lat at Local Noon	57°N	48°S	10°N	43°N	45°N
Dates on Orbit	2-9 Dec 92	3-11 Feb 95	7-18 Sep 95	12-20 Nov 95	7-19 Aug 97
Limb View Hours	30	50	120	50	>300
Sunlit L. V. Hours	25	30	70	40	200
Fractional moon	0.56-0.99	0.17-0.86	0.97-0.31	0.73-0.04	0.25-1.0
A _p Index	2-12	2-30	3-39	3-10	5-16
F10.7 solar flux (10 ⁻²² W/cm ² Hz)	112-116	79-84	68-72	71-75	76-84
Tangent Height km	80-330	80-400	80-400	80-400	80-300
Spectrograph Slit	Parallel	--- Parallel Night, Vertical Day ---			
Data Focus	Terminator	Near Global	Equatorial	Near Global	Near Global




Figure 1. Orbit tracks showing latitudinal extent and relation to terminator for the five Space Shuttle missions.




Figure 2. GLO-5 and GLO-6 being integrated onto the IEH2 Hitchhiker bridge in preparation for the STS-85 flight.

Data analysis efforts included the development of the SIBYL code to manipulate GLO data, evaluation of thruster-atmosphere interactions, identification of atmospheric morphology, and validation/verification of atmospheric compensation codes. The latter work tested codes which were developed by the Air Force Research Laboratory and its contractors, using Modtran4 atmospheric simulations to remove the effects of atmospheric components and scattering from the measured radiance in hyperspectral data. The resulting reflectance spectral cubes are then available for use as inputs to AFRL data categorization routines.

Ground-based observation of the ionosphere was also performed during RF-heater excitation experiments. A prototype dual-spectrograph system was assembled to demonstrate the importance of full visible range spectral observations during ionospheric excitation experiments, such as those planned for the Air Force HAARP site in Gakona, AK. The prototype was fielded during the January 1998 RF-heater campaign at Arecibo, PR, and returned positive identification of molecular bands during ionospheric excitation events. These molecular bands had not been previously predicted by the ionospheric heating community.

Publications resulting from this work:

1. A. L. Broadfoot, B. R. Sandel, D. J. Knecht, R. Viereck, E. Murad, *A Panchromatic Spectrograph with Supporting Monochromatic Imagers*, Applied Optics, **31**, 3083 (1992).
2. A. L. Broadfoot, E. Anderson, P. Sherard, D. J. Knecht, R. A. Viereck, C. P. Pike, E. Murad, J. E. Elgin, L. S. Bernstein, I. L. Kofsky, D. L. A. Rall, J. Blaha, F. L. Culbertson, *Spectrographic Observation at Wavelengths Near 630 nm of the Interaction Between the Atmosphere and the Space Shuttle Exhaust*, J. Geophys. Res., **97**, 19,501 (1992).
3. J. E. Johnston and A. L. Broadfoot, *Midlatitude Observations of the Night Airglow: Implications to Quenching Near the Mesopause*, J. Geophys. Res., **98**, 21,593 (1993).
4. J. A. Gardner, R. A. Viereck, E. Murad, S. T. Lai, D. J. Knecht, C. P. Pike, A. L. Broadfoot, W. J. McNeil, *Mg⁺ and Other Metallic Emissions Observed in the Thermosphere*, Proc. SPIE, **2266**, 242 (1994).
5. J. E. Johnston, D. B. Hatfield, A. L. Broadfoot, *Synthetic spectra for the Arizona Airglow Experiment*, Proc. SPIE, **2266**, 480 (1994).
6. J. A. Gardner, R. A. Viereck, E. Murad, D. J. Knecht, C. P. Pike, A. L. Broadfoot, E. R. Anderson, *Simultaneous Observations of Neutral and Ionic Magnesium in the Thermosphere*, Geophys. Res. Lett., **22**, 2119 (1995).
7. R. A. Viereck, E. Murad, D. J. Knecht, C. P. Pike, L. S. Bernstein, J. B. Elgin, A. L. Broadfoot, *The interaction of the atmosphere with the space shuttle thruster plume: The NH (A-X) 336-nm emission*, J. Geophys. Res., **101**, 5371-5380 (1996).
8. J. A. Gardner, E. Murad, D. J. Knecht, R. A. Viereck, C. P. Pike, A. L. Broadfoot, *Thermospheric Metal Emissions: Update on GLO Mg⁺ Measurements*, Proc. SPIE, **2830**, 64 (1996).
9. J. A. Gardner, E. Murad, D. J. Knecht, R. A. Viereck, C. P. Pike, A. L. Broadfoot, *Shuttle-based Measurements: GLO Ultraviolet Earthlimb View*, Proc. SPIE, **2831**, 125 (1996).
10. R. A. Viereck, E. Murad, S. T. Lai, D. J. Knecht, C. P. Pike, J. A. Gardner, A. L. Broadfoot, E. R. Anderson, W. J. McNeil, *Mg⁺ and other metallic emissions observed in the thermosphere*, Adv. Space Res., **18**, 61 (1996).
11. A. L. Broadfoot, D. B. Hatfield, E. R. Anderson, T. C. Stone, B. R. Sandel, J. A. Gardner, E. Murad, D. J. Knecht, C. P. Pike, R. A. Viereck, *The N₂ Triplet Band Systems in the Dayglow*, J. Geophys. Res., **102**, 11576 (1997).
12. D. J. Knecht, E. Murad, R. A. Viereck, C. P. Pike, A. L. Broadfoot, E. R. Anderson, D. B. Hatfield, T. C. Stone, B. R. Sandel, *Arizona Airglow Experiment as flown on Four Space-Shuttle Missions*, Advances in Space Research, **19**, 627-630 (1997).
13. A.L. Broadfoot, *Remote Sensing from the Space Shuttle and Space Stations*, Adv. Space Res., **19**, 623-626 (1997).

14. J. A. Gardner, E. Murad, R. A. Viereck, D. J. Knecht, C. P. Pike, A. L. Broadfoot, *GLO Observations of E- and F-Region Metal Atoms and Ions*, Adv. Space Res., **21**, 867 (1998).
15. J. A. Gardner and A. L. Broadfoot, *Meteor Metals in the Ionosphere: GLO Shuttle-Based Observations*, Proc. AIAA 99-0505, (1999).
16. J. A. Gardner, E. Murad, W. J. McNeil, S. T. Lai, A. L. Broadfoot, *Analysis and Modeling of the GLO-1 Observations of Meteoric Metals in the Thermosphere*, J. Atmospheric and Solar-Terrestrial Physics, **61**, 545 (1999).
17. F. T. Djuth, P. A. Bernhardt, C. A. Tepley, J. A. Gardner, M. C. Kelley, A. L. Broadfoot, L. M. Kagan, M. P. Sulzer, J. H. Elder, C. Selcher, B. Isham, C. Brown, H. C. Carlson, *Large Airglow Enhancements Produced via Wave-Plasma Interactions in Sporadic-E*, Geophys. Res. Lett., **26**, 1557-1560 (1999).
18. J. A. Gardner, G. W. Felde, E. Meidunas, M. Hoke, J. Chetwynd, G. P. Anderson, S. J. Lipson, P. S. Armstrong, B. Pukall, J. B. Mello, L. S. Jeong, S. M. Adler-Golden, A. Berk, L. S. Bernstein, P. K. Acharya, M. W. Matthew, *Considerations In Atmospheric Compensation Of Spectral Image Data*, Proc. SPIE, **3756**, 363 (1999).
19. G. P. Anderson, A. Berk, L. S. Bernstein, J. Chetwynd, P. K. Acharya, H. Dothe, M. W. Matthew, S. M. Adler-Golden, R. J. Ratkowski, G. W. Felde, J. A. Gardner, M. L. Hoke, S. C. Richtsmeier, B. Pukall, J. Mello, L. S. Jeong, *MODTRAN4: Radiative Transfer Modeling for Remote Sensing and Atmospheric Correction*, Proc. SPIE, **3866-02** (1999).
20. A. L. Broadfoot and P. J. Bellaire, Jr., *Bridging the Gap Between Ground-Based and Space-Based Observations of the Night Airglow*, J. Geophys. Res., **104**, 17127-17138 (1999).
21. A. L. Broadfoot and T. C. Stone, *$N_2 + O^+$ charge-exchange reaction and the dayglow N_2^+ emission*, J. Geophys. Res., **104**, 17145-17155 (1999).

STS-85

THE ARIZONA AIRGLOW EXPERIMENT

GLO-5 AND GLO-6

END-OF-MISSION REPORT

SUBMITTED BY: Lyle Broadfoot, P.I.
University of Arizona

The science operations of the GLO-5 and GLO-6 experiments on board STS-85 have been a spectacular success! The two GLO instruments performed spectral and imagery measurements on several optical data targets:

Target	# orbits	Comment
ionospheric dayglow	73	43 in "preferred" attitude
ionospheric night airglow	68	all in good attitudes
Aurora Australis	12	some ground-to-deep-space scans
planets, stars (calibration data)	20	pointing/detector response checks
Perseid meteor shower metals	68	data set in night airglow
Hale-Bopp comet	5	
cometary OH	3	
space vehicle glow	2	
OMS engine firings	2	2 events at altitude change
"Near-stereo" viewing	20	1 at 90° others at 40° angle
SEH, CRISTA coordinated scans	30	

Both GLO instruments performed flawlessly, with better-than-expected thermal responses and no serious anomalies. GLO observations occurred in all of the shuttle's routine attitudes, and a total of 25 gigabytes of ITM (Ionosphere, Thermosphere, Mesosphere) data were recorded,

rivaling the total for GLO-2, -3, and -4 on STS-63, -69, and -74 in 1995. Lessons learned from, and instrument upgrades since, those flights significantly improved both the operation of and scientific return from the instruments.

The GLO instruments are essentially identical imaging spectrographs. Each contains a spectrograph suite covering 120 - 930 nm in several steps, using Intensified Charge Coupled Device (ICCD) detectors. Detector chips are chosen for best sensitivity in each of several wavelength regions. The GLOs also include filtered imagers that are dedicated to monitoring OH, O₂, and O in the airglow. A low-light-level video camera is on-board to assist in pointing the instrument, and all optics are co-aligned and boresighted to a single view-center. On GLO-6, an EUV diode is also mounted to support the SEH solar-viewing experiment.

Dayglow and airglow measurements were performed in two modes: (i) spectrograph slit perpendicular to the earth limb, and (ii) slit near-parallel to the earth limb. The first mode was accessed during -YLV, -XLV, and +XLV attitudes. With the long axis of the spectrograph slit perpendicular to the limb, the GLOs record altitude-resolved optical data. For dayglow measurements, the two GLOs operated with their Fields-Of View (FOV) stacked, to provide continuous coverage from about 100 to 300 km tangent height at the earth limb. For night airglow, one GLO was positioned with its FOV parallel to the limb to provide maximum sensitivity with a narrow height range. The other GLO had its FOV tilted to provide about 40 km of viewing height at the limb, to provide simultaneous coverage of the 70 to 110 km region in which the important airglow features are found. The GLOs each collected 43 orbits of dayglow and 38 orbits of airglow in this mode.

The slit-nearly-parallel mode is the only mode accessible during -ZLV (bay-to-earth) attitudes. Here, the 15° wedges that tilt the scan platforms forward allowed GLO to maintain the 40 km range in tangent heights during the day. Without the wedges, no useful data would have been collected during Day periods in the bay-to-earth attitudes. At night, airglow measurements proceeded as described above for the other attitudes. Each GLO collected 30 orbits of dayglow and 30 orbits of airglow in the bay-to-earth attitudes.

The Southern Aurora produced a superb target during STS-85. On about 8 separate orbits, intense auroral activity was measured from 100 - 300 km altitudes using the normal airglow method. During 4 of the inertial orbit attitudes, the aurora was measured as the earth

limb swept through the FOV. On these orbits, the aurora were measured to their upper limits at several hundred kilometers altitudes.

GLO also performed measurements on stars to provide calibration data. Because stars are distant point sources, they are used to verify the co-alignment of each GLO's 12 detectors. Standard stars with well-known spectra, including Vega, Spica, Sirius, and Beta Centauri, are measured by the GLO instruments to provide spectral calibration of the instruments. The many inertial shuttle attitudes provided an excellent range of opportunities to calibrate the GLO instruments and scan platforms.

Because of the fortuitous launch on August 7, *Discovery* was on orbit during the Perseid meteor shower. This annual event lasts for several days, and August 12 is the peak of the ground-observable "shooting stars." Although meteor showers are a small portion of the total mass loading of the upper atmosphere by particles from space, the showers represent intense periods where the daily mass loading greatly exceeds the average. The Perseid meteor shower is expected to increase temporarily the natural metal layers near 90 km altitude by as much as 10 - 20%. The GLO instruments were able to scan this region before, during, and after the shower peak and will report on the average sodium layer density over the course of the mission.

The Hale-Bopp comet is a widely recognized target of opportunity for STS-85. As did several other experiments, the GLO instruments obtained spectra at Hale-Bopp's position during at least 4 orbits. The GLOs also attempted to measure electronically excited hydroxyl (OH) molecules that would be required by-products of "small comets". Post-flight data analysis will determine the scientific value of these data sets.

The GLO program historically has been interested in measuring thruster firings and effluents, as well as looking for the space vehicle glow phenomenon. During STS-85, the orbit adjust burns on Flight Day 11 were observed at multiple angles, looking especially for optical signatures of thruster gases downstream of the burns. Shuttle glow observation experiments were performed during several eclipse periods in the bay-to-earth configuration. GLOs also performed measurements in support of SEH and CRISTA: a solar Extreme UltraViolet diode on GLO-6 monitored solar output, and both GLOs supported CRISTA "Sprites" and thermospheric experiments. For each of these types of experiments, the data sets are small and will require post-mission analysis prior to commenting on their scientific value.

The GLOs also performed the definitive "near-stereo" viewing experiment. On prior GLO flights, only one instrument flew at a time. Although the scientific return from those earlier flights has been outstanding, the ionospheric data has always lacked an important dimension. The earlier data has shown altitude profiles along the shuttle flight track, yielding a 2-dimensional view of any selected spectral feature. With both GLOs on the same flight, we extend these measurements to include the depth dimension by pointing the instruments to view tangent points from different angles with only a short time delay between the measurements. The views were set to nearly 90° separation and viewed common points at 3 minute time delays. This data set will yield 3-dimensional (spatial) data on important atmospheric dynamics, such as the magnesium ion lines near 280 nm, which can be used as tracers for the earth's electric field.

The flight of GLO-5 and GLO-6 on STS-85 has been an absolute success! Both machines were completely nominal in performance. Ionospheric data included over 70 orbits of ionospheric data, dozens of stellar calibration points, and experiments focused on Jupiter, Hale-Bopp, cometary OH, shuttle glow, thruster firings, and coordinated observations with other STS-85 payloads. A key improvement to the instruments was the addition of a filter to remove second-order effects inside the ultraviolet spectrographs. This filter blocked interference from the oxygen 130.4 nm harmonic at 260.8 nm and allowed for GLOs' first positive identification of the strong iron ion emission line at 261.4 nm. Coupled with the magnesium ion lines at 280 nm and the sodium D lines at 589 nm, the iron data will greatly improve the understanding of metal cycling in the ionosphere. All results discussed here are from "quick-look" reduction of quality-control sampled data from each GLO. Far less than 1% of the total data set is available now, with the remainder being eagerly awaited.